

A Guide to Managing Potential Side Effects

Following the suggestions in this guide may help you feel better during treatment with RASONQUE

INDICATION

RASONQUE is a prescription medicine used to treat adults with a type of pancreatic cancer called pancreatic adenocarcinoma:

- that has spread to other parts of the body, **and**
- who have received at least 1 prior treatment or who cannot receive a combination of cancer treatments.

It is not known if RASONQUE is safe and effective in children.

Please see Important Safety Information on [pages 10–11](#) and full [Prescribing Information](#), including [Medication Guide](#).

 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg

How to use this guide

This guide can help you learn about the possible side effects of RASONQUE and how to manage them. Ask your care team to explain anything you don't understand about managing side effects.

If you have a side effect, always let your care team know right away. In addition, ask a loved one or caregiver to help you track and manage side effects.

Additional materials that may be helpful are available on the Resources page at RASONQUE.com. You can use the Treatment Journal to record side effects and take notes. The Care Team Discussion Guide can help you during conversations at your appointments.

This guide has two sections.

Identifying side effects: Includes a list of common side effects and what they look like

Managing side effects: Explains how you and your care team may be able to manage side effects so you can feel better

Use the menu at the bottom of each page to quickly jump between sections.

Visit RASONQUE.com for more helpful resources.



Actor portrayal.

The content here is for informational and educational purposes only and is not intended for medical advice. It does not take the place of medical advice or treatment from a physician. You should always consult your own doctor or a qualified healthcare professional for specific health concerns and questions.

Please see Important Safety Information on pages 10–11 and full [Prescribing Information](#), including [Medication Guide](#).

 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg

Identifying potential side effects

Review what some common side effects from treatment with RASONQUE may look like



Rash

Red, dry skin and/or small bumps that commonly occur on your neck, back, and face



Diarrhea

More than your usual number of stools. They can be loose or watery



Swelling or sores in the mouth (stomatitis)

Inflammation or irritation in the mouth



Upset stomach (nausea and/or vomiting)

Feeling queasy like you might throw up and/or throwing up the food and liquid in your stomach

Other common side effects include: tiredness or weakness, stomach (abdominal) pain, swelling in the arms and legs, decreased appetite, and bleeding.

Please see Important Safety Information on pages 10–11 and full Prescribing Information, including Medication Guide.



RASONQUE[™]
(daraxonrasib) tablets 100 mg
150 mg

Identifying potential side effects (cont)

Your doctor may adjust your dose if you have side effects with RASONQUE

- **97.1% of people** who received RASONQUE did not have to stop treatment because of side effects
- People who received RASONQUE may have had their dose paused or adjusted to help manage side effects
 - 69% of people had their dose paused
 - 37% of people had their dose reduced

RASONQUE can cause serious side effects, including: skin reactions, swelling or sores in the mouth (stomatitis), diarrhea, tears in your stomach or intestines (gastrointestinal perforation), and lung or breathing problems (pneumonitis). RASONQUE can harm your unborn baby.

Partner with your care team

- Talk to your care team about how you're feeling. They can help you better understand any side effects you have
- Keep your care team's contact information handy. Save it in your phone or write it down somewhere convenient
- Keep track of any side effects you have. If you have a rash, take a photo that you can share with your care team

Please see Important Safety Information on pages 10–11 and full Prescribing Information, including Medication Guide.



 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg

Managing potential side effects

Always talk to your care team about any side effects you have right away. They may be able to help you manage them so you can feel better.

There are steps you and your caregiver can take to help manage certain side effects on your own at home.

Your doctor may also pause or adjust your dose of RASONQUE. This may help reduce symptoms of side effects.



Actor portrayal.

Remember to let your care team know right away about any side effects you have, even if they are not bothering you that much

Please see Important Safety Information on pages 10–11 and full [Prescribing Information](#), including [Medication Guide](#).



 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg

5

Guidelines for management

Follow the guidelines in this section to help prevent or manage some common side effects of RASONQUE. Remember to always tell your care team right away about any side effects you have as soon as possible.



Guidelines to help prevent and manage rash

To help prevent and manage rash, you **SHOULD**:

- ✓ Apply certain medications to your skin (face and body) before your first dose; talk to your care team
- ✓ Use a scent-free and soothing moisturizer to keep your skin hydrated
- ✓ Limit sun exposure and wear protective clothing, including a hat, if you are in the sun
- ✓ Use sunscreen that is SPF 30 or higher
 - Apply at least 30 minutes before going out into the sun. Reapply every 2 hours
- ✓ Discuss taking oral antibiotics with your care team
 - Your doctor may recommend you start taking these before your treatment with RASONQUE starts
 - If you have questions or concerns about taking antibiotics, talk with your care team
- ✓ Keep your skin clean; consider using an over-the-counter antiseptic spray (eg, hypochlorous acid)
- ✓ Use warm (not hot) water while bathing

Always talk to your care team about how you're feeling. They can help you better understand any side effects you have. They may be able to help you manage them so you can feel better.

Please see Important Safety Information on pages 10–11 and full Prescribing Information, including Medication Guide.



 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg

Guidelines for management (cont)

Your care team can help you manage rash

Rash can be mild, moderate, or severe. If you develop a rash, take a photo of it to share with your care team. This way, they can help you monitor it and decide what action, if any, is needed.

If your rash is more severe (bleeding or painful), your care team may have additional directions for how to manage it.

Contact your care team at the first sign of a rash, even if it isn't bothering you



TIP: Download the Rash Management Card from the Resources page of RASONQUE.com for more information.



Please see Important Safety Information on pages 10–11 and full Prescribing Information, including Medication Guide.



RASONQUE[™]
(daraxonrasib) tablets 100 mg
150 mg

7

Guidelines for management (cont)

Potential side effect	DO	DO NOT
 Diarrhea	✓ Contact your care team if you have diarrhea (or more stools than usual), if your diarrhea gets worse, or if it does not improve ✓ Follow your care team's instructions on when and how to take any medicine they recommend ✓ Stay hydrated by drinking plenty of water ✓ Try changing your diet until the diarrhea improves – One option is to eat a BRAT diet (bananas, rice, applesauce, toast)	✗ Drink sugary beverages
 Swelling or sores in the mouth (stomatitis)	✓ Contact your care team if you develop mouth sores ✓ Keep up with your oral hygiene routine, including brushing and flossing your teeth at least twice a day ✓ Rinse your mouth with saline (salt water) – You can make your own rinse solution at home by mixing 1 teaspoon of salt and 1 teaspoon of baking soda with 4 cups of water. Only use 1 mouthful at a time to rinse	✗ Eat rough and sharp-edged foods, like potato chips ✗ Use tobacco or alcohol (including beverages and mouthwashes) ✗ Eat foods with extreme (hot or cold) temperatures

Please see Important Safety Information on pages 10–11 and full Prescribing Information, including Medication Guide.



RASONQUE[™]
(daraxonrasib) tablets 100 mg
150 mg

Guidelines for management (cont)

Potential side effect	DO	DO NOT
 Upset stomach (nausea and/or vomiting)	✓ Contact your care team if you have an upset stomach ✓ Follow your care team’s instructions on when and how to take any medicine they recommend ✓ Eat small, frequent meals ✓ Sip water or clear liquids throughout the day	✗ Eat overly fatty, greasy, or spicy foods



TIP: Download the Treatment Journal from the Resources page of RASONQUE.com to help track any side effects you have. This information can help your care team decide how to move forward.

Please see Important Safety Information on pages 10–11 and full Prescribing Information, including Medication Guide.



 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg

INDICATION

RASONQUE is a prescription medicine used to treat adults with a type of pancreatic cancer called pancreatic adenocarcinoma:

- that has spread to other parts of the body, **and**
- who have received at least 1 prior treatment or who cannot receive a combination of cancer treatments.

It is not known if RASONQUE is safe and effective in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about RASONQUE?

RASONQUE can cause serious side effects, including:

- **Skin reactions.** Skin reactions may be severe. To help reduce your risk of skin reactions and protect your skin when starting and during treatment with RASONQUE:

- use a topical steroid cream on your face and chest
- use a moisturizing cream on your skin
- limit your time in the sun and use sunscreen (broad spectrum SPF 30 or higher) and other sun protection measures such as clothing and hats if you must be in the sun
- take any other medicines if prescribed by your healthcare provider

Tell your healthcare provider right away if you develop any signs or symptoms of skin reactions, including:

- rash
- red, swollen, and painful skin
- dry skin
- itching
- around your finger or toenails
- cracked skin

- **Swelling or sores in the mouth (stomatitis).** Mouth sores may be severe. Your healthcare provider may prescribe a mouthwash or other medicines to treat your mouth reactions. Tell your healthcare provider right away if you develop trouble eating, talking, swallowing, or develop any new or worsening signs or symptoms in your mouth, including:

- pain
- redness
- swelling
- ulcers or sores

- **Diarrhea.** Diarrhea may be severe. If you develop diarrhea during treatment with RASONQUE, your healthcare provider may prescribe medicines to treat your diarrhea. Tell your healthcare provider right away if you develop new or worsening diarrhea or if the number of bowel movements you have in a day increases by 6 or more.

- **Tears in your stomach or intestines (gastrointestinal perforation).** Gastrointestinal perforation may be severe and lead to death. Tell your healthcare provider right away if you develop any of the following signs or symptoms:

- tenderness or pain in your stomach area (abdomen) that is severe or does not go away
- blood in the stool or black stool that looks like tar
- nausea, vomiting, or vomiting blood
- fever or chills

Please see additional Important Safety Information on [page 11](#) and full [Prescribing Information](#), including [Medication Guide](#).



RASONQUE[™]
(daraxonrasib) tablets 100 mg
150 mg

10

IMPORTANT SAFETY INFORMATION (cont)

- **Lung or breathing problems (pneumonitis).** Lung or breathing problems may be severe and lead to death. Tell your healthcare provider right away if you develop any new or worsening breathing problems, including:

- cough - shortness of breath - trouble breathing - wheezing

The most common side effects of RASONQUE include:

- rash
- diarrhea
- swelling or sores in the mouth
- nausea
- tiredness or weakness
- vomiting
- stomach (abdominal) pain
- swelling in the arms and legs
- decreased appetite
- bleeding

Certain abnormal blood tests are common during treatment with RASONQUE. Your healthcare provider will monitor you for abnormal blood tests and treat you if needed.

Your healthcare provider may decrease your dose, or temporarily or permanently stop treatment with RASONQUE if you develop certain side effects.

Before taking RASONQUE, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant. RASONQUE can harm your unborn baby.

Females who are able to become pregnant:

- Your healthcare provider will do a pregnancy test before you start treatment with RASONQUE.
- Use effective birth control (contraception) during treatment with RASONQUE and for 1 week after the last dose.
- Tell your healthcare provider right away if you become pregnant during treatment with RASONQUE.

Males with female partners who are able to become pregnant:

- Use effective contraception during treatment with RASONQUE and for 1 week after the last dose.
- are breastfeeding or plan to breastfeed. It is not known if RASONQUE passes into your breast milk. **Do not** breastfeed during treatment and for 1 week after your last dose of RASONQUE.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. RASONQUE can affect the way other medicines work, and other medicines can affect how RASONQUE works, and may increase your risk of side effects.

These are not all the possible side effects of RASONQUE. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or www.fda.gov/Safety/MedWatch.

Please see additional Important Safety Information on page 10 and full Prescribing Information, including Medication Guide.



 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg

11

Managing side effects is key to your overall care

Contacting your care team can make all the difference

Need help starting the conversation with your care team?

Download the Care Team Discussion Guide from the Resources page of RASONQUE.com.

Visit RASONQUE.com for more helpful resources.

Revisit this guide while taking RASONQUE to help manage side effects

INDICATION

RASONQUE is a prescription medicine used to treat adults with a type of pancreatic cancer called pancreatic adenocarcinoma:

- that has spread to other parts of the body, **and**
- who have received at least 1 prior treatment or who cannot receive a combination of cancer treatments.

It is not known if RASONQUE is safe and effective in children.

Please see Important Safety Information on pages 10–11 and full Prescribing Information, including Medication Guide.

You are encouraged to report negative side effects of Revolution Medicines prescription drugs by calling 1-844-2-REVMED (1-844-273-8633). If you prefer to report these to the FDA, call 1-800-FDA-1088.



This product information is intended for US consumers only.
RASONQUE is a trademark of Revolution Medicines, Inc.
© 2026 Revolution Medicines, Inc. RVMD-USA-00386(v1.0) 08/26

 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg

12