

**FIRST
APPROVED
TREATMENT OF ITS KIND**
ORAL, NON-CHEMOTHERAPY

MORE TIME for what matters

RASONQUE is for adults (18 and older) with a type of **PANCREATIC CANCER** called pancreatic adenocarcinoma that has spread to other parts of the body, and who have received at least 1 prior treatment or who cannot receive a combination of cancer treatments.

[Table of contents >](#)

In a clinical trial, people who were treated with RASONQUE had a median overall survival of 13.2 months vs 6.6 months for those treated with chemotherapy. Median overall survival is the length of time when half the people on treatment are still living.

Actor portrayal.
RASONQUE is not for use in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about RASONQUE?

RASONQUE can cause serious side effects, including:

- **Skin reactions.** Skin reactions may be severe. To help reduce your risk of skin reactions and protect your skin when starting and during treatment with RASONQUE:
 - use a topical steroid cream on your face and chest
 - use a moisturizing cream on your skin
 - limit your time in the sun and use sunscreen (broad spectrum SPF 30 or higher) and other sun protection measures such as clothing and hats if you must be in the sun
 - take any other medicines if prescribed by your healthcare provider

Please see additional Important Safety Information on pages 19–20 and full Prescribing Information, including Medication Guide.

 **RASONQUE™**
(daraxonrasib) tablets 100 mg
150 mg

Get the answers you need when starting RASONQUE

Use the table of contents below to find information and resources to help support you during treatment.


<u>Your Guide to RASONQUE</u>	3
<u>How does RASONQUE work?</u>	4
<u>How was RASONQUE studied?</u>	6
<u>What results did people have with RASONQUE?</u>	8
<u>What are some of the possible side effects of RASONQUE?</u>	12
<u>What should I do if I have side effects?</u>	13
<u>How do I take RASONQUE?</u>	15
<u>How will I receive RASONQUE?</u>	17
<u>Where can I find RASONQUE support services?</u>	18
<u>Important Safety Information</u>	19

Visit [RASONQUE.com](https://www.rasonque.com) for additional support and resources.

Please see Important Safety Information on pages 19–20 and full Prescribing Information, including [Medication Guide](#).

 **RASONQUE**™
(daraxonrasib) tablets 100 mg
150 mg

Your guide to RASONQUE



Actor portrayal.
RASONQUE is not for use in children.

Starting a targeted treatment can bring new possibilities.

RASONQUE is used to treat adults:

- who have pancreatic cancer that has spread to other parts of the body (metastatic), and
- who have received at least 1 prior treatment or who cannot receive a combination of cancer treatments

RASONQUE is not chemotherapy. RASONQUE is a targeted treatment that's taken orally, once a day.

Read on to learn more about targeted treatment with RASONQUE

IMPORTANT SAFETY INFORMATION (cont)

What is the most important information I should know about RASONQUE? (cont)

RASONQUE can cause serious side effects, including: (cont)

• Skin reactions. (cont)

Tell your healthcare provider right away if you develop any signs or symptoms of skin reactions, including:

- rash
- red, swollen, and painful skin around your finger or toenails
- dry skin
- cracked skin

Please see additional Important Safety Information on [pages 19–20](#) and full [Prescribing Information](#), including [Medication Guide](#).



 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg

How does RASONQUE work?

RASONQUE helps stop or slow down the growth of cancer cells by targeting RAS



RAS is found in cells in your body. It sends signals that help cells know when to start and stop growing. Sometimes, these signals can cause cells to grow out of control. This can lead to cancer.

RASONQUE is a targeted treatment. It looks for RAS that is sending signals for cells to grow. When RASONQUE finds RAS, it stops these signals. This helps **stop or slow down the growth of cancer cells.**

RASONQUE was studied in people whose cancer did or did not have a RAS mutation

IMPORTANT SAFETY INFORMATION (cont)

What is the most important information I should know about RASONQUE? (cont)

RASONQUE can cause serious side effects, including: (cont)

- **Swelling or sores in the mouth (stomatitis).** Mouth sores may be severe. Your healthcare provider may prescribe a mouthwash or other medicines to treat your mouth reactions. Tell your healthcare provider right away if you develop trouble eating, talking, swallowing, or develop any new or worsening signs or symptoms in your mouth, including:
 - pain
 - swelling
 - redness
 - ulcers or sores


Please see additional Important Safety Information on [pages 19–20](#) and full Prescribing Information, including [Medication Guide](#).



RASONQUE™
(daraxonrasib) tablets 100 mg
150 mg


RASONQUE is a targeted treatment

Targeted treatments work differently than chemotherapy



Chemotherapy

kills or slows the growth of cells that are growing quickly, including cancer cells and healthy cells.



Targeted treatments

seek out and attack specific cells that cause cancer to grow. Because they look for these cells, they are less likely to affect healthy cells.



Healthy cells



Cancer cell (tumor)

IMPORTANT SAFETY INFORMATION (cont)

What is the most important information I should know about RASONQUE? (cont)

RASONQUE can cause serious side effects, including: (cont)

- **Diarrhea.** Diarrhea may be severe. If you develop diarrhea during treatment with RASONQUE, your healthcare provider may prescribe medicines to treat your diarrhea. Tell your healthcare provider right away if you develop new or worsening diarrhea or if the number of bowel movements you have in a day increases by 6 or more.

Please see additional Important Safety Information on [pages 19–20](#) and full Prescribing Information, including [Medication Guide](#).



RASONQUE[™]
(daraxonrasib) tablets 100 mg
150 mg


How was RASONQUE studied?

RASONQUE was studied in a clinical trial called RASolute 302

The trial included 500 adults:

- ✓ With pancreatic cancer that had spread to other parts of the body (metastatic)
- ✓ Whose cancer progressed after prior chemotherapy
- ✓ Whose cancer did or did not have a RAS mutation*

People received either:



*The trial included 459 people whose cancer had a RAS G12 mutation and 41 people whose cancer did not have a RAS G12 mutation.

†The chemotherapy that people received was chosen by their doctor from the following combinations: modified FOLFIRINOX (leucovorin + fluorouracil + irinotecan + oxaliplatin); gemcitabine + nab-paclitaxel; FOLFOX (leucovorin + fluorouracil + oxaliplatin); or liposomal irinotecan + fluorouracil/leucovorin.

Please see Important Safety Information on [pages 19–20](#) and full Prescribing Information, including [Medication Guide](#).



 **RASONQUE™**
(daraxonrasib) tablets 100 mg / 150 mg

The trial compared RASONQUE to chemotherapy

The main goals of the trial were to study:

- **Overall survival:** How long people lived after starting treatment
- **Progression-free survival:** How long people lived without their cancer growing or spreading after starting treatment

The trial also studied:

- **Tumor response:** How many people had their tumors shrink
- **Duration of response:** How long a person's cancer responded to treatment
- **Safety:** What side effects people had after starting treatment
- **Health-related quality of life:** How long people lived without their health-related quality of life getting worse

Please see Important Safety Information on [pages 19–20](#) and full [Prescribing Information](#), including [Medication Guide](#).



 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg

What results did people have with RASONQUE?

RASONQUE was studied in people whose metastatic pancreatic cancer had a RAS G12 mutation and those whose cancer did not have a RAS G12 mutation. Researchers looked at the trial data from people whose cancer had a RAS G12 mutation first. Then, they looked at the data from everyone in the trial.

Of people whose cancer had a RAS G12 mutation:

- The median overall survival for people who received RASONQUE was **13.2 months** compared to **6.6 months** for those who received chemotherapy
- The median progression-free survival for people who received RASONQUE was **7.3 months** compared to **3.5 months** for those who received chemotherapy

Of everyone in the trial:

RASONQUE helped people live longer compared to chemotherapy



- People who received RASONQUE had a **60% reduction in their risk of death** compared to those who received chemotherapy

Median overall survival is the length of time when half the people on treatment are still living.

IMPORTANT SAFETY INFORMATION (cont)

What is the most important information I should know about RASONQUE? (cont)

RASONQUE can cause serious side effects, including: (cont)

- **Tears in your stomach or intestines (gastrointestinal perforation).** Gastrointestinal perforation may be severe and lead to death. Tell your healthcare provider right away if you develop any of the following signs or symptoms:
 - tenderness or pain in your stomach area (abdomen) that is severe or does not go away
 - nausea, vomiting, or vomiting blood
 - blood in the stool or black stool that looks like tar
 - fever or chills

Please see additional Important Safety Information on [pages 19–20](#) and full [Prescribing Information](#), including [Medication Guide](#).



RASONQUE™
(daraxonrasib) tablets 100 mg
150 mg

What results did people have with RASONQUE?

Of everyone in the trial:

RASONQUE doubled the time people lived without cancer growing or spreading compared to chemotherapy



51% reduction in their risk of death or of cancer growing or spreading for people who received RASONQUE compared to those who received chemotherapy

Median progression-free survival is the length of time when half the people in the trial are still living without their cancer growing or spreading.

IMPORTANT SAFETY INFORMATION (cont)

What is the most important information I should know about RASONQUE? (cont)

RASONQUE can cause serious side effects, including: (cont)

- **Lung or breathing problems (pneumonitis).** Lung or breathing problems may be severe and lead to death. Tell your healthcare provider right away if you develop any new or worsening breathing problems, including:
 - cough
 - trouble breathing
 - shortness of breath
 - wheezing


Please see additional Important Safety Information on [pages 19–20](#) and full Prescribing Information, including [Medication Guide](#).



RASONQUE™
(daraxonrasib) tablets 100 mg
150 mg

Additional results

Tumor response



People who had
their tumors shrink*

For those who
received RASONQUE

30%
of 248 people

For those who
received chemotherapy

11%
of 252 people



People who had
their tumors stabilize†

48.1%
of 237 people

38.6%
of 241 people

Stabilize (stable disease) means that changes in tumor sizes were too small to be reported as growing or shrinking. This may be due to how tumors behave naturally rather than any effect of the medicine. Please consider that these results are not definitive, and individual results may vary.

These results were not tested for statistical significance and not designed to show differences between treatments. Statistical significance is a mathematical measure that shows whether a difference between groups is greater than what might be expected (predicted) to happen by chance alone.

*Evaluated in everyone in the trial.

† Evaluated in people who had measurable disease before starting treatment in the trial.

IMPORTANT SAFETY INFORMATION (cont)

What is the most important information I should know about RASONQUE? (cont)

The most common side effects of RASONQUE include:

- rash
- diarrhea
- swelling or sores in the mouth
- nausea
- tiredness or weakness
- vomiting
- stomach (abdominal) pain
- swelling in the arms and legs
- decreased appetite
- bleeding

Certain abnormal blood tests are common during treatment with RASONQUE. Your healthcare provider will monitor you for abnormal blood tests and treat you if needed. Your healthcare provider may decrease your dose, or temporarily or permanently stop treatment with RASONQUE if you develop certain side effects.

Please see additional Important Safety Information on [pages 19–20](#) and full Prescribing Information, including [Medication Guide](#).



RASONQUE™
(daraxonrasib) tablets 100 mg
150 mg

Additional results

Health-related quality of life

Health-related quality of life is how a person feels about their overall physical and mental health over a period of time.

How was health-related quality of life studied?

People in the trial completed questionnaires about their health-related quality of life each month. They answered each question with a numeric score. This helped researchers find out how much time people had before certain measures got worse. These measures included overall quality of life/health and pain.

- Consider these results carefully. The results shown here may be unrelated to treatment. In addition, the trial was an open-label study. This means that both the researchers and the participants knew which treatment they were receiving

	For those who received RASONQUE	For those who received chemotherapy
Median time without pain getting worse	9.2 months	3.8 months
Median time without overall quality of life/health getting worse	5.7 months	2.6 months

The surveys used were the European Organization for Research and Treatment of Cancer Quality of Life Questionnaire (EORTC QLQ-C30) and EORTC QLQ-PAN 26.

Median is the middle number in a set of numbers.

IMPORTANT SAFETY INFORMATION (cont)

What is the most important information I should know about RASONQUE? (cont)

Before taking RASONQUE, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant. RASONQUE can harm your unborn baby.

Females who are able to become pregnant:

- Your healthcare provider will do a pregnancy test before you start treatment with RASONQUE.
- Use effective birth control (contraception) during treatment with RASONQUE and for 1 week after the last dose.
- Tell your healthcare provider right away if you become pregnant during treatment with RASONQUE.

Males with female partners who are able to become pregnant:

- Use effective contraception during treatment with RASONQUE and for 1 week after the last dose.

Please see additional Important Safety Information on [pages 19–20](#) and full Prescribing Information, including [Medication Guide](#).



 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg

What are some of the possible side effects of RASONQUE?

Serious common side effects were:



Rash



Diarrhea



Swelling or sores
in the mouth

Additional common side effects were:

- Nausea
- Tiredness or weakness
- Vomiting
- Stomach (abdominal) pain
- Swelling in the arms and legs
- Decreased appetite
- Bleeding

RASONQUE can cause serious side effects, including: skin reactions, swelling or sores in the mouth (stomatitis), diarrhea, tears in your stomach or intestines (gastrointestinal perforation), and lung or breathing problems (pneumonitis). RASONQUE can harm your unborn baby.

These are not all the possible side effects of RASONQUE. Call your doctor for medical advice about side effects.

Talk to your care team right away if you experience any side effects

Please see Important Safety Information on [pages 19–20](#) and full [Prescribing Information](#), including [Medication Guide](#).



 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg

What should I do if I have side effects?

Call your care team right away

Everyone on your care team has the same goal: to help you manage side effects and continue treatment.



Your care team may include:

- Oncologists
- Nurse navigators
- Physician assistants
- Pharmacists
- Nurses and nurse practitioners
- Dietitian/nutritionist

You and your caregiver are the leaders of your team! Make sure to use your voice and connect with your care team when you need to.

Your care team will recommend certain medicines and products that may reduce side effects

Please see Important Safety Information on [pages 19–20](#) and full [Prescribing Information](#), including [Medication Guide](#).



RASONQUE[™]
(daraxonrasib) tablets 100 mg
150 mg

Rash is the most common side effect of RASONQUE

- 87% of people who received RASONQUE had a rash
- Your care team will recommend certain medicines or products to start before taking RASONQUE to help reduce the risk of rash

Visit [RASONQUE.com](https://www.rasonque.com) for tips on managing rash

Your doctor may adjust your dose if you have side effects with RASONQUE


- 97.1% of people who received RASONQUE did not have to stop treatment because of side effects
- People who received RASONQUE may have had their dose paused or adjusted to help manage side effects
 - 69% of people had their dose paused
 - 37% of people had their dose reduced

Please see Important Safety Information on [pages 19–20](#) and full [Prescribing Information](#), including [Medication Guide](#).



 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg

How do I take RASONQUE?



Once daily

300 mg as two 150-mg tablets

Tablets not actual size.


**RASONQUE is an oral treatment
that can fit into your daily routine**



**At the same time
each day**



With or without food



At home or on the go

- Take RASONQUE exactly as your doctor tells you to take it. Do not change your dose or stop taking RASONQUE unless your doctor tells you to
- Swallow RASONQUE tablets whole. **Do not** chew, crush, or split tablets

Your doctor may adjust your dose if you have side effects with RASONQUE. Tell your doctor about all the medicines you take, and do not start any new medicines without talking to your doctor.

Please see Important Safety Information on [pages 19–20](#) and full [Prescribing Information](#), including [Medication Guide](#).



 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg

Table of
contents

About
RASONQUE

Study
& results

Side
effects

Taking
RASONQUE

Support
services

Important Safety
Information

Helpful tips to keep your treatment routine on track



Set an alarm or ask a friend/loved one to remind you



Use a calendar or pill organizer



Take RASONQUE at the same time you do another daily activity, like brushing your teeth



Consider having a snack to help with nausea

- If you miss a dose, take the dose as soon as you remember. If it has been more than 4 hours, skip the missed dose and take your next dose at your next regularly scheduled time. **Do not** take 2 doses at the same time to make up for a missed dose
- If you vomit after taking a dose, do not take an extra dose. Take your next dose at your regularly scheduled time

Please see Important Safety Information on [pages 19–20](#) and full Prescribing Information, including [Medication Guide](#).



 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg

How will I receive RASONQUE?

Your RASONQUE prescription will be filled by a specialty pharmacy

Specialty pharmacies handle specialty medicines

- A specialty pharmacy offers more services than the retail pharmacies you may be more familiar with
- Pharmacists with oncology experience are available to help answer questions about side effects and your dosing schedule

Your doctor's office and specialty pharmacy will work together to deliver RASONQUE to you on time, by mail

- Once your oncologist sets up your prescription for RASONQUE, a specialty pharmacist will reach out
- Be sure to answer their call or return their voicemail message, as it may be from a number you do not recognize
- Your specialty pharmacy will be sure to deliver RASONQUE to your door at the address you provide

Please see Important Safety Information on [pages 19–20](#) and full [Prescribing Information](#), including [Medication Guide](#).



 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg



Support that moves you forward

Working with your provider to enroll in (ON)Path means hands-on support from a dedicated (ON)Path Patient Access Advocate who will connect you or your loved one with personalized programs and resources to help start treatment as soon as possible.

Subject to terms and conditions, you may be eligible for:



(ON)Path Patient
Access Advocates



Quick Start Program



Commercial Copay
Program



Patient Assistance
Program

Have questions?
We're (ON) it.

Call an (ON)Path Patient Access Advocate at **1-844-ONPATH0**
(844-667-2840), Monday-Friday, 8 AM to 8 PM EST

INDICATION

RASONQUE is a prescription medicine used to treat adults with a type of pancreatic cancer called pancreatic adenocarcinoma:

- that has spread to other parts of the body, **and**
- who have received at least 1 prior treatment or who cannot receive a combination of cancer treatments.

It is not known if RASONQUE is safe and effective in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about RASONQUE?

RASONQUE can cause serious side effects, including:

- **Skin reactions.** Skin reactions may be severe. To help reduce your risk of skin reactions and protect your skin when starting and during treatment with RASONQUE:
 - use a topical steroid cream on your face and chest
 - use a moisturizing cream on your skin
 - limit your time in the sun and use sunscreen (broad spectrum SPF 30 or higher) and other sun protection measures such as clothing and hats if you must be in the sun
 - take any other medicines if prescribed by your healthcare provider

Tell your healthcare provider right away if you develop any signs or symptoms of skin reactions, including:

 - rash
 - itching
 - red, swollen, and painful skin around your finger or toenails
 - dry skin
 - cracked skin
- **Swelling or sores in the mouth (stomatitis).** Mouth sores may be severe. Your healthcare provider may prescribe a mouthwash or other medicines to treat your mouth reactions. Tell your healthcare provider right away if you develop trouble eating, talking, swallowing, or develop any new or worsening signs or symptoms in your mouth, including:
 - pain - redness - swelling - ulcers or sores

- **Diarrhea.** Diarrhea may be severe. If you develop diarrhea during treatment with RASONQUE, your healthcare provider may prescribe medicines to treat your diarrhea. Tell your healthcare provider right away if you develop new or worsening diarrhea or if the number of bowel movements you have in a day increases by 6 or more.
- **Tears in your stomach or intestines (gastrointestinal perforation).** Gastrointestinal perforation may be severe and lead to death. Tell your healthcare provider right away if you develop any of the following signs or symptoms:
 - tenderness or pain in your stomach area (abdomen) that is severe or does not go away
 - nausea, vomiting, or vomiting blood
 - blood in the stool or black stool that looks like tar
 - fever or chills
- **Lung or breathing problems (pneumonitis).** Lung or breathing problems may be severe and lead to death. Tell your healthcare provider right away if you develop any new or worsening breathing problems, including:
 - cough
 - shortness of breath
 - trouble breathing
 - wheezing

The most common side effects of RASONQUE include:

- | | |
|----------------------------------|---------------------------------|
| • rash | • vomiting |
| • diarrhea | • stomach (abdominal) pain |
| • swelling or sores in the mouth | • swelling in the arms and legs |
| • nausea | • decreased appetite |
| • tiredness or weakness | • bleeding |

Certain abnormal blood tests are common during treatment with RASONQUE. Your healthcare provider will monitor you for abnormal blood tests and treat you if needed.

Please see additional Important Safety Information on page 20 and full Prescribing Information, including Medication Guide.

 **RASONQUE™**
(daraxonrasib) tablets 100 mg
150 mg

IMPORTANT SAFETY INFORMATION (cont)

Your healthcare provider may decrease your dose, or temporarily or permanently stop treatment with RASONQUE if you develop certain side effects.

Before taking RASONQUE, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant. RASONQUE can harm your unborn baby.

Females who are able to become pregnant:

- Your healthcare provider will do a pregnancy test before you start treatment with RASONQUE.
- Use effective birth control (contraception) during treatment with RASONQUE and for 1 week after the last dose.
- Tell your healthcare provider right away if you become pregnant during treatment with RASONQUE.

Males with female partners who are able to become pregnant:

- Use effective contraception during treatment with RASONQUE and for 1 week after the last dose.
- are breastfeeding or plan to breastfeed. It is not known if RASONQUE passes into your breast milk. **Do not** breastfeed during treatment and for 1 week after your last dose of RASONQUE.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. RASONQUE can affect the way other medicines work, and other medicines can affect how RASONQUE works, and may increase your risk of side effects.

These are not all the possible side effects of RASONQUE. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or www.fda.gov/Safety/MedWatch.

Please see additional Important Safety Information on page 19 and full Prescribing Information, including Medication Guide.

 **RASONQUE**™
(daraxonrasib) tablets 100 mg
150 mg

An oral, non-chemotherapy treatment for adults with metastatic pancreatic cancer who have received at least 1 prior treatment or who cannot receive a combination of cancer treatments



Actor portrayal.
RASONQUE is not for use in children.

Embrace more everyday moments

Find more support and resources during your treatment with RASONQUE

Visit RASONQUE.com to learn more.

IMPORTANT SAFETY INFORMATION (cont)

Before taking RASONQUE, tell your healthcare provider about all of your medical conditions, including if you: (cont)

- are breastfeeding or plan to breastfeed. It is not known if RASONQUE passes into your breast milk. **Do not** breastfeed during treatment and for 1 week after your last dose of RASONQUE.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. RASONQUE can affect the way other medicines work, and other medicines can affect how RASONQUE works, and may increase your risk of side effects.

These are not all the possible side effects of RASONQUE. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or www.fda.gov/Safety/MedWatch.

Please see additional Important Safety Information on pages 19–20 and full Prescribing Information, including Medication Guide.

You are encouraged to report negative side effects of Revolution Medicines prescription drugs by calling 1-844-2-REVMED (1-844-273-8633). If you prefer to report these to the FDA, call 1-800-FDA-1088.



This product information is intended for US consumers only.
RASONQUE and (ON)Path are trademarks of Revolution Medicines, Inc.
© 2026 Revolution Medicines, Inc. RVMD-USA-00385(v1.0) 08/26

 **RASONQUE**[™]
(daraxonrasib) tablets 100 mg
150 mg

Table of
contents

About
RASONQUE

Study
& results

Side
effects

Taking
RASONQUE

Support
services

Important Safety
Information